

Negative-Parity Baryons in Quenched Anisotropic Lattice QCD

Y. Nemoto^{a*}, N. Nakajima^b, H. Matsufuru^c and H. Suganuma^d

^aRIKEN-BNL Research Center, Brookhaven National Laboratory, Upton 11973, USA

^bCenter of Medical Information Science, Kochi Medical School, Kochi 783-8505, Japan

^cYukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

^dFaculty of Science, Tokyo Institute of Technology, Tokyo 152-8551, Japan

We study negative-parity baryon spectra in quenched anisotropic lattice QCD. The negative-parity baryons are measured as the parity partner of the ground-state baryons. In addition to the flavor octet and decuplet baryons, we pay much attention to the flavor-singlet negative-parity baryon as a three-quark state and compare it with the $\Lambda(1405)$ baryon. Numerical results of the flavor octet and decuplet negative-parity baryon masses are close to experimental values of lowest-lying negative-parity baryons, while the flavor-singlet baryon is much heavier than $\Lambda(1405)$. This indicates that the $\Lambda(1405)$ would be a multi-quark state such as the $N\bar{K}$ molecule rather than the flavor-singlet 3 quark state.

1. INTRODUCTION

In the lattice QCD Monte Carlo calculation, the spectroscopy of the ground-state hadrons in the quenched approximation has been almost established and reproduces the experimental values within 10% deviations. On the other hand, the spectroscopy of the excited-state baryons is a current interesting subject in lattice QCD. In this study, we focus on the low-lying negative-parity baryons, especially the parity partner of the ground-state baryons and the flavor-singlet baryon, $\Lambda(1405)$. $\Lambda(1405)$ is a mysterious baryon in the quark model viewpoint. It is the lightest negative-parity baryon though it has strangeness. Our purpose of this paper is to clarify whether a naive three valence quark picture can explain the mass of $\Lambda(1405)$ using the characteristics of quenched lattice QCD, and to investigate spectra of the parity partners of the ground-state baryons of the flavor octet and decuplet states[1, 2]. The other possible candidate for the $\Lambda(1405)$ state is the $N\bar{K}$ (5-quark) state. It is useful to employ the quenched approximation to know whether the dominant component of $\Lambda(1405)$ is the three-quark state, because the overlap with the three-quark and five-quark states is absent and hence the properties of the purely three-quark state can be singled out.

*Y.N. was supported by the center-of-excellence (COE) program at YITP, Kyoto University in most stage of this work and thanks RIKEN, Brookhaven National Laboratory and the U.S. Department of Energy for providing the facilities essential for the completion of this work.

Table 1

Baryon operators. C is the charge conjugate matrix.

Octet	$B_\alpha(\Sigma^0) = (C\gamma_5)_{\beta\gamma}[u_\alpha(d_\beta s_\gamma - s_\beta d_\gamma) - d_\alpha(s_\beta u_\gamma - u_\beta s_\gamma)]$
Octet (Λ)	$B_\alpha(\Lambda) = (C\gamma_5)_{\beta\gamma}[u_\alpha(d_\beta s_\gamma - s_\beta d_\gamma) + d_\alpha(s_\beta u_\gamma - u_\beta s_\gamma) - 2s_\alpha(u_\beta d_\gamma - d_\beta u_\gamma)]$
Decuplet	$B_{\alpha k}(\Sigma^{*0}) = (C\gamma_k)_{\beta\gamma}[u_\alpha(d_\beta s_\gamma + s_\beta d_\gamma) + d_\alpha(s_\beta u_\gamma + u_\beta s_\gamma) + s_\alpha(u_\beta d_\gamma + d_\beta u_\gamma)]$
Singlet	$B_\alpha(\Lambda_1) = (C\gamma_5)_{\beta\gamma}[u_\alpha(d_\beta s_\gamma - s_\beta d_\gamma) + d_\alpha(s_\beta u_\gamma - u_\beta s_\gamma) + s_\alpha(u_\beta d_\gamma - d_\beta u_\gamma)]$

2. LATTICE SETUP

We employ the standard Wilson gauge action and the $O(a)$ improved Wilson quark action with tadpole improvement. We adopt the anisotropic lattice because high resolution in the temporal direction makes us easy to follow the change of correlation of heavy baryons. We take the renormalized anisotropy as $\xi = a_\sigma/a_\tau = 4$, where a_σ and a_τ are the spatial and temporal lattice spacings, respectively. The simulation is performed on three lattices for which the parameters are well tuned and the errors are rather well evaluated in Ref.[5]. The sizes of the lattices are $12^3 \times 96$ ($\beta = 5.75, a_\sigma^{-1} = 1.34(6)$ GeV), $16^3 \times 128$ ($\beta = 5.95, a_\sigma^{-1} = 1.099(9)$ GeV) and $20^3 \times 160$ ($\beta = 6.10, a_\sigma^{-1} = 1.871(14)$ GeV). The scale a_σ^{-1} is determined from the K^* meson mass. Determination of the parameters is described in Ref.[5].

On the quark mass, we adopt four different values, which roughly cover around the strange quark mass. We use the standard baryon operator listed in Table 1 which survive in the nonrelativistic limit. The negative-parity baryons can be also measured by these operators using the parity projection. They are nothing but the parity partners of the ground-state baryons. In the source operator, each quark is smeared with the Gaussian function of the width ~ 0.4 fm.

3. RESULTS AND DISCUSSION

The lattice data and the fit results of the baryon for the $\beta = 6.10$ lattice are shown in Fig. 1. The physical u, d and s quark masses are determined with the π and K meson masses. We fit the mass data to the linear form for the pseudoscalar meson mass square.

In order to compare our lattice results with experimental values, various baryon masses for $\beta = 6.10$ lattice together with the experimental values are shown in Fig. 2. As for the negative-parity baryons, most of the present lattice results comparatively well reproduces the experimental spectra of the lowest-lying negative-parity baryons, in spite of relatively large statistical errors. (For the light baryons such as the nucleon and the delta, it is reported that there are non-analytic terms in the chiral extrapolation of the nucleon in the quenched level, which may cause the nucleon and the delta heavier than the experimentally observed masses to some extent.) However, the lattice result of the flavor-singlet negative-parity baryon is much heavier than the experimentally observed $\Lambda(1405)$. Note here that the flavor-singlet baryon has one strange valence quark and therefore the ambiguity from the chiral extrapolation should be less than that of the nucleon and the delta. Nevertheless, the discrepancy between the lattice result and experimental value of $\Lambda(1405)$ is the largest of all. Even if we take the quenching effect of 10% level into account, this large discrepancy cannot be understood. Therefore, the numerical result for




Figure 1. Spectra of the positive- and negative-parity baryons for the $\beta = 6.10$ lattice. Octet and decuplet baryons are shown in the left figure and octet(Λ) and singlet baryons in the right figure. The open symbols denote lattice data and the filled symbols the results fitted from the linear form of the chiral extrapolation.

the flavor-singlet baryon exhibits a different feature from other channels, which indicates that it is unacceptably heavier than $\Lambda(1405)$. Thus, we conclude that the present lattice result indicates that the three valence quark picture fails to represent the experimentally observed $\Lambda(1405)$, i.e., the overlap with the 3Q component of $\Lambda(1405)$ is small. The other possible candidate for $\Lambda(1405)$ is a five-quark $N\bar{K}$ state. Lattice calculations of the $N\bar{K}$ (5Q) state will also elucidate the nature of $\Lambda(1405)$.

We focus on other negative-parity baryons. From Fig. 2, we see that the lattice results of the flavor octet and decuplet sectors are all close to the observed lowest-lying negative parity baryons, $N(1535)$, $\Lambda(1670)$, $\Sigma(1620)$, $\Xi(1690)$ and $\Delta(1700)$, although they have a relatively large statistical error. The identification of the parity partner of the sigma baryon is interesting because the evidence of existence of $\Sigma(1620)$ is only fair[6]. As for the Ξ baryons, although the Ξ with $J^P = \frac{1}{2}^-$ baryon is not known experimentally, we can identify the parity partner as $\Xi(1690)$ from our results. For the flavor decuplet baryons, $\Delta(1700)$ can be regarded as the parity partner of $\Delta(1232)$.

4. CONCLUSION

We have studied the mass spectra of the negative-parity baryons, which are the parity partners of the ground-state (positive-parity) baryons, and the flavor-singlet baryons in the quenched anisotropic lattice QCD. The positive- and negative-parity baryon masses are extracted from the same baryon correlators with the parity projection.

Compared with the experimental values, our present results of the lattice calculation show that the possible assignment of the parity partners of the ground-state baryons for octet and decuplet is the lowest-lying negative-parity baryons. It is also interesting to relate our lattice data to some phenomenological models for parity doublet baryons[4]




Figure 2. Various baryon masses from the $\beta = 6.10$ lattice. For the negative-parity baryons, we add the experimental values of $N(1535)$, $\Lambda(1670)$, $\Sigma(1620)$, $\Xi(1690)$, $\Delta(1700)$ and $\Lambda(1405)$.

for future.

On the other hand, the mass of the negative-parity flavor-singlet baryon is much heavier than that of $\Lambda(1405)$, whose behavior is quite different from other negative-parity baryons. This large discrepancy between the lattice result and experimental spectrum seems to imply that the three valence quark picture fails to represent the $\Lambda(1405)$ state, i.e., the overlap with the three-quark state of $\Lambda(1405)$ is small. For more definite understanding of $\Lambda(1405)$ spectrum, one needs more extensive work about the flavor-singlet baryon such as the observation of the $N\bar{K}$ (five-quark) state. Such a study is interesting even at the quenched level, where the dynamical quark-loop effect including the quark-antiquark pair creation is absent and then the quark-level constitution of hadrons is definitely clear in the simulation.

REFERENCES

1. Y. Nemoto, N. Nakajima, H. Matsufuru and H. Suganuma, Phys. Rev. D **68** (2003) 094505.
2. N. Nakajima, H. Matsufuru, Y. Nemoto and H. Suganuma, AIP **CP594** (2001) 349; H. Suganuma, N. Ishii, H. Matsufuru, Y. Nemoto, T. T. Takahashi, **CP644** (2002) 366.
3. S. Sasaki, et al., Phys. Rev. D **65** (2002) 074503; M. Göckeler et al., Phys. Lett. B **532** (2002) 63; W. Melnitchouk et al., Phys. Rev. D **67** (2003) 114506 F.X. Lee et al., Nucl. Phys. Proc. Suppl. **119** (2003) 296 ; UKQCD Collaboration, C.M. Maynard and LHP Collaboration, D.G. Richards, Nucl. Phys. Proc. Suppl. **119** (2003) 287.
4. Y. Nemoto et al., Phys. Rev. D **57** (1997) 4124; D. Jido et al., Nucl. Phys. A **671** (2000) 471; D. Jido et al., Prog. Theor. Phys. **106** (2001) 873.
5. H. Matsufuru, T. Onogi and T. Umeda, Phys. Rev. D **64** (2001) 114503.
6. Particle Data Group, Phys. Rev. D **66** (2002) 1.